

# Investigation of Hamstring Tendon Graft Fixation for the Reconstruction of Anterior Cruciate Ligament using Interference Screw Merely or in Combination with Supplementary Staple: A Clinical Trial

## Abstract

**Background:** Hamstring graft tendon for anterior cruciate ligament (ACL) reconstruction is a common approach worldwide. Tibial side graft fixation to achieve appropriate stability is a serious concern, worldwide. The current study aims to compare the outcomes of mere use of interference screw for fixation of hamstring tendon graft versus the use of interference screw plus supplementary staple. **Materials and Methods:** This is a randomized clinical trial conducted on 53 patients who underwent ACL reconstruction from 2016 to 2018. The study population was randomly divided into two groups: graft fixation with interference screw only and interference screw plus supplementary staple. Postoperative recovery time, postoperative clinical examinations, and the scale of the International Knee Documentation Committee were assessed for participants and compared between two groups. **Results:** Comparison of two groups regarding demographic information, including age, gender distribution, postoperative recovery time, and body mass index, showed no statistical difference ( $P > 0.05$ ). Postoperative Pivot test was insignificantly positive in 4 (16.7%) cases of screw interference with supplementary staple while it was positive in 3 (10.7%) cases with screw interference only approach ( $P = 0.98$ ). IKCD index was not statistically different between two groups postoperatively ( $P = 0.72$ ), while IKCD scores changed significantly following the surgical procedure, regardless of the type of the surgical procedure ( $P < 0.001$ ). **Conclusion:** Use of supplementary staple beside interference screw was as successful as mere use of interference screw for fixation of hamstring tendon autologous graft of the ACL reconstruction, regarding force withstanding. The comparison of the two approaches revealed no remarkable difference.

**Keywords:** Anterior cruciate ligament, anterior cruciate ligament reconstruction, bone screw, bone-patellar tendon-bone grafts, sutures

## Introduction

Reconstruction of anterior cruciate ligament (ACL) is among the most common orthopedic procedures, which could successfully restore the stability and function of the injured knee following the ligament rupture and allow the person to return to daily routine activities.<sup>[1,2]</sup>

However, the graft choice for achieving optimal reconstruction outcomes has been a great challenge worldwide; two graft options including autologous central third patellar tendon with the associated bone (bone-patellar tendon-bone) (BTB) and hamstring grafts consisting of semitendinosus and gracilis are the most popular ones.<sup>[3,4]</sup> Reports in the literature have presented equivalent success rate of these two approaches, while the use of

hamstring graft leads to less morbidity and greater strength of the graft, making this approach of remarkable favor by orthopedists.<sup>[5,6]</sup>

Soft tissue only grafts without a bone plug such as the approach of hamstring tendons graft are considered as the weakest point for initial ACL reconstruction. This fact occurs due to the forces against the graft that are mostly parallel to the tibia tunnel, while the tibial metaphysis is a relatively softer bone than the femoral tunnel.<sup>[7,8]</sup>

Another remarkable point about such procedures that involve soft tissue grafts only without bone integration is the considerable required time for its integration occurrence, unlike the bone-to-bone integrations.<sup>[9]</sup>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [WKHLRPMedknow\\_reprints@wolterskluwer.com](mailto:WKHLRPMedknow_reprints@wolterskluwer.com)

**How to cite this article:** Mousavi SH, Masoumi O, Akbariaghdam H, Mohammadsharifi G. Investigation of hamstring tendon graft fixation for the reconstruction of anterior cruciate ligament using interference screw merely or in combination with supplementary staple: A clinical trial. *Adv Biomed Res* 2020;9:52.

Sayed Hamid Mousavi,  
Omid Masoumi<sup>1</sup>,  
Hossein Akbariaghdam,  
Ghasem Mohammadsharifi<sup>1</sup>

Department of Orthopedics,  
Kashani Hospital, Isfahan  
University of Medical Sciences,  
<sup>1</sup>Department of Orthopedics,  
Al-Zahra Hospital, Isfahan  
University of Medical Sciences,  
Isfahan, Iran

**Address for correspondence:**  
Dr. Omid Masoumi,  
Department of Orthopedics,  
Al-Zahra Hospital, Isfahan  
University of Medical  
Sciences, Isfahan, Iran.  
E-mail: [omidmasoumy@gmail.com](mailto:omidmasoumy@gmail.com)

**Received:** 06 December 2019  
**Revised:** 21 January 2020  
**Accepted:** 18 May 2020  
**Published:** 30 September 2020

### Access this article online

**Website:** [www.advbiores.net](http://www.advbiores.net)

**DOI:** 10.4103/abr.abr\_257\_19

### Quick Response Code:








Figure 1: Consort diagram of the studied population

This scale was assessed before the surgical procedure and then within at least 8 months following the surgical procedure.<sup>[14]</sup>

The tunnel view radiography including anterior–posterior, upright, and 35° knee flexion view was taken preoperatively and at the end of the physiotherapy sessions as the follow-up study. The radiographies were performed to assess the severity of the injury and compartment degeneration.

### Study analysis

Obtained data were entered into SPSS-20 (IBM; Chicago; The United States) and analyzed. Descriptive data were presented in mean and percentages, while for analytics, Chi-square test, *t*-test, and Fisher’s exact test were used. *P* < 0.05 was considered statistically significant.

### Results

In the current study, eligibility of 60 patients was assessed,

among which four did not meet the inclusion criteria and 56 were randomly allocated to the interventions in equal groups of screw interference with supplementary staple ( $n = 28$ ) and screw interference only ( $n = 28$ ). All of the operated cases in the screw interference with supplementary staple finished the study protocol, while four in the latter group lost to refer for follow-up visit [Figure 1].

The mean age and mean BMI of the study population were  $30.58 \pm 6.38$  years and  $27.12 \pm 4.02$  kg/m<sup>2</sup>, with a predominance of the male gender, with 66.03% of the distribution. Most of the study population presented right-sided involvement of ACL (67.9%).

Comparison of two groups regarding demographic information including age, gender distribution, postoperative recovery time, time of postoperative follow-up visit, and BMI showed no statistical differences ( $P > 0.05$ ). Postoperative ADT and Lachman test were negative for both of the groups, while Pivot test was positive in 4 (16.7%) cases of screw interference with supplementary staple and in 3 (10.7%) cases with screw interference the only approach. Comparison of the two groups revealed no statistical difference ( $P = 0.98$ ). Table 1 represents the mentioned information in details.

The clinical knee examinations, including Lachman, Pivot, and ADTs, have been examined before the surgical procedure and then compared with the assessments performed following it. Table 2 demonstrates the status of these tests before and after the procedure. Based on the findings of Table 2, both groups represented a significant improvement in functional examinations, following whether the screw interference only or in combination with a staple ( $P < 0.05$ ). Postoperative comparison of two groups

regarding Pivot test positivity revealed no remarkable difference between those treated with the screw interference with a supplementary staple and those who were merely treated with screw interference approach ( $P = 0.98$ ).

IKCD index assessments are demonstrated in Table 3. Based on Table 3, two groups were not statistically different before the surgical procedures ( $P = 0.69$ ). Besides, comparison of two groups, postoperatively, showed no statistical difference ( $P = 0.72$ ), while IKCD scores changed significantly following the surgical procedure regardless of the type of the surgical procedure ( $P < 0.001$ ).

## Discussion

Although hamstring autograft has become of great interest for the reconstruction of ACL, a notifying concern about its inability to tolerate the initial postoperative forces of rehabilitation made scientists to raise novel approaches.<sup>[10,15]</sup>

The primary principals of this study were to compare the outcomes of the interference screw only versus interference screw with supplementary staple among the patients who had undergone ACL reconstruction using hamstring tendon autograft. As we conducted our study on two groups of the patients similar in age, gender distribution, BMI, and the side of injured ACL, the probable role of these factors on the conclusion of our study was eliminated. Postoperative examination of two groups showed no statistical differences regarding clinical laxity tests of Pivot shift, Lachman, and ADT. Besides, patients' subjective declarations of IKDC score were not statistically different between the two groups. In general, we found no remarkable benefit of supplementary staple use beside interference screw.

**Table 1: Comparison of demographics and postoperative examination tests between two groups of study population**

Variable	Screw interference with supplementary staple ( $n=24$ )	Screw interference only ( $n=28$ )	P	Test
Gender* (male/female)	15 (62.5)/9 (37.5)	20 (71.4)/8 (28.6)	0.19	Chi-square
Age**	31.81±6.37	29.83±6.33	0.22	T-test
Body mass index**	27.13±3.88	27.10±4.22	0.97	T-test
Involved side* (right/left)	16 (66.7)/8 (33.3)	20 (71.4)/8 (28.6)	0.29	T-test
Postoperative recovery time**	6.53±2.01	7.23±1.43	0.12	T-test
Postoperative follow-up time (month)**	2.20±9.80	2.36±10.30	0.40	T-test

\*n (%), \*\*Mean±SD. SD: Standard deviation

**Table 2: Comparison of clinical examinations of the knee before the surgical procedure and after that considering the type of the surgery**

Variable	Group	Before, n (%)	After, n (%)	P*
Lachman	Screw interference with a supplementary staple (+/-)	24 (100)/0 (0)	0 (0)/24 (100)	<0.001
	Screw interference only (+/-)	28 (100)/0 (0)	0 (0)/28 (100)	<0.001
Pivot	Screw interference with supplementary staple (+/-)	24 (100)/0 (0)	4 (16.7)/20 (83.3)	<0.001
	Screw interference only (+/-)	28 (100)/0 (0)	3 (10.7)/25 (89.3)	<0.001
Anterior drawer test	Screw interference with supplementary staple (+/-)	24 (100)/0 (0)	0 (0)/24 (100)	<0.001
	Screw interference only (+/-)	28 (100)/0 (0)	0 (0)/28 (100)	<0.001

\*McNemar t-test

**Table 3: Comparison of the International Knee Documentation Committee scale between two groups of the study population**

Surgical technique	n	Mean±SD		P*	P* Difference means
		Before	After		
Screw interference with supplementary staple	24	59.67±19.83	79.47±14.41	<0.001	0.80
Screw interference only	28	61.57±16.98	80.70±12.82	<0.001	

\*Paired *t*-test, \*\**t*-test. SD: Standard deviation

Reports in the literature have presented acceptable outcomes of eccentric interference screw fixation through the ACL reconstruction using hamstring tendon, regarding the stability of reconstructed ligament for withstanding against the forces,<sup>[8,16]</sup> while some more recent studies have presented that the use of supplementary staple beside screw has remarkable beneficial in comparison to the mere use of the screws.<sup>[13,17]</sup> This theory is still a great matter of research as further evaluations showed that even if the use of staple would provide better stability against forces, this device may pose considerable complications such as pain and reoperation requirement.<sup>[1,14]</sup>

Teo *et al.* conducted a similar study as they evaluated their patients with similar means including Lachman and Pivot tests, and IKCD, and presented consistent outcomes as they found no difference between the mere interference screw and interference screw plus staple, neither in clinical examinations nor through subjective IKCD assessments. The superiority of their study was its longer duration of follow-up than us, which presented staple-related complications, including postoperative pain due to kneeling and reoperation requirement for device removal.<sup>[14]</sup>

Hill *et al.* performed another study on females who underwent ACL reconstruction with autograft of hamstring tendon using whether interference screw fixation or in combination with a supplementary staple. Although, in a similar declaration to us, they presented no significant difference between two groups regarding postoperative IKCD, they declared that treated patients with supplementary staple plus metal screws presented notably reduced laxity of their knee through clinical Lachman examinations as compared to the mere use of metal screws. They interpreted that the density of proximal tibial metaphysis bone may have decreased, following the injury and further manipulations.<sup>[13]</sup>

Bauer *et al.* also compared the outcomes of screw interference only or in combination with staple. Although they assessed the benefits of the methods based on the cross-sectional area of the reconstructed ligament, yield load, failure point on 10 mm and device failure, similar to our outcomes, they found no superiority for the staple use added to screw interference.<sup>[18]</sup>

Although we have not found any failure in our study, our patients were approximately followed only for a year while most of the studies have presented that the ultimate tendon to bone integration would occur within 2 years following the reconstruction procedure.<sup>[2]</sup> Nevertheless, there are reports representing shorter time required for achieving the desired integration as Pinczewski *et al.* represented that only 15 weeks is required to achieve the expected microscopic and macroscopic hamstring autograft–bone interface.<sup>[19]</sup> On the other hand, animal models have demonstrated that signs of tendon-to-bone integration can be detected within 4 months.<sup>[20]</sup> These findings were confirmed by Pinczewski *et al.* who presented both microscopic and macroscopic hamstring tendon autograft integration within 15 postoperative weeks.<sup>[19]</sup> The other study by Rodeo *et al.* presented the biological interface of bone to the reconstructed tendon within 8 weeks of the operation.<sup>[21]</sup> Therefore, it seems that our study protocol for postoperative follow-up was adequately presented by other authors as well.<sup>[22]</sup>

Of limitations of our study is not to assess the procedure-related complications, while the symptoms due to kneeling have made the theory on the selection of patients for staple use added to interference screw. Because pain experience poses significant bothersome to the patient, it poses a burden of secondary surgery procedure for its removal.<sup>[14]</sup>

## Conclusion

Use of supplementary staple beside interference screw was as successful as mere use of interference screw for fixation of hamstring tendon autologous graft of ACL reconstruction regarding force withstanding. Comparison of two approaches revealed no remarkable difference.

## Limitations

The most significant limitation of the current study is its little number of studied populations. Therefore, further studies with larger sample populations are recommended.



## Acknowledgment

We are grateful to Kashani Hospital's head office, nurses, and employees.

## Financial support and sponsorship

The study was sponsored by Isfahan University of Medical Sciences.

## Conflicts of interest

There are no conflicts of interest.

## References

1. Pinczewski LA, Lyman J, Salmon LJ, Russell VJ, Roe J, Linklater J. A 10-year comparison of anterior cruciate ligament reconstructions with hamstring tendon and patellar tendon autograft: A controlled, prospective trial. *Am J Sports Med* 2007;35:564-74.
2. De Wall M, Scholes CJ, Patel S, Coolican MR, Parker DA. Tibial fixation in anterior cruciate ligament reconstruction: A prospective randomized study comparing metal interference screw and staples with a centrally placed polyethylene screw and sheath. *Am J Sports Med* 2011;39:1858-64.
3. Goldblatt JP, Fitzsimmons SE, Balk E, Richmond JC. Reconstruction of the anterior cruciate ligament: Meta-analysis of patellar tendon versus hamstring tendon autograft. *Arthroscopy* 2005;21:791-803.
4. Poolman RW, Farrokhyar F, Bhandari M. Cumulative meta-analysis and clinically relevant sensitivity analysis show evidence for hamstring tendon autograft superiority compared to bone patellar-tendon bone autograft in ACL reconstruction. *Moving towards Evidence-Based Orthopaedic Surgery*.2007;12:256-262.
5. Wagner M, Kääh MJ, Schallock J, Haas NP, Weiler A. Hamstring tendon versus patellar tendon anterior cruciate ligament reconstruction using biodegradable interference fit fixation: A prospective matched-group analysis. *Am J Sports Med* 2005;33:1327-36.
6. Li S, Chen Y, Lin Z, Cui W, Zhao J, Su W. A systematic review of randomized controlled clinical trials comparing hamstring autografts versus bone-patellar tendon-bone autografts for the reconstruction of the anterior cruciate ligament. *Arch Orthop Trauma Surg* 2012;132:1287-97.
7. Duncan RM, Meuffels E, Docter PT, van Dongen RA, Kleinrensink GJ, Jan AN, *et al.* Novel insights into anterior cruciate ligament injury. *Erasmus University Rotterdam* 2010;26:65.
8. Scranon PE Jr., Bagenstose JE, Lantz BA, Friedman MJ, Khalfayan EE, Auld MK. Quadruple hamstring anterior cruciate ligament reconstruction: A multicenter study. *Arthroscopy* 2002;18:715-24.
9. Bartz RL, Mossoni K, Tyber J, Tokish J, Gall K, Siparsky PN. A biomechanical comparison of initial fixation strength of 3 different methods of anterior cruciate ligament soft tissue graft tibial fixation: Resistance to monotonic and cyclic loading. *Am J Sports Med* 2007;35:949-54.
10. Kousa P, Järvinen TL, Vihavainen M, Kannus P, Järvinen M. The fixation strength of six hamstring tendon graft fixation devices in anterior cruciate ligament reconstruction. Part I: Femoral site. *Am J Sports Med* 2003;31:174-81.
11. Brand J Jr., Weiler A, Caborn DN, Brown CH Jr., Johnson DL. Graft fixation in cruciate ligament reconstruction. *Am J Sports Med* 2000;28:761-74.
12. Gwynne-Jones DP, Draffin J, Vane AG, Craig RA, McMahon SF. Failure strengths of concentric and eccentric implants for hamstring graft fixation. *ANZ J Surg* 2008;78:177-81.
13. Hill PF, Russell VJ, Salmon LJ, Pinczewski LA. The influence of supplementary tibial fixation on laxity measurements after anterior cruciate ligament reconstruction with hamstring tendons in female patients. *Am J Sports Med* 2005;33:94-101.
14. Teo WW, Yeoh CS, Wee TH. Tibial fixation in anterior cruciate ligament reconstruction: Is supplementary staple fixation necessary? *J Orthopaed Surg* 2017;25:2309499017699743.
15. Ibrahim SA, Al-Kussary IM, Al-Misfer AR, Al-Mutairi HQ, Ghafar SA, El Noor TA. Clinical evaluation of arthroscopically assisted anterior cruciate ligament reconstruction: Patellar tendon versus gracilis and semitendinosus autograft. *Arthroscopy* 2005;21:412-7.
16. Laxdal G, Kartus J, Eriksson BI, Faxén E, Sernert N, Karlsson J. Biodegradable and metallic interference screws in anterior cruciate ligament reconstruction surgery using hamstring tendon grafts: Prospective randomized study of radiographic results and clinical outcome. *Am J Sports Med* 2006;34:1574-80.
17. Tetsumura S, Fujita A, Nakajima M, Abe M. Biomechanical comparison of different fixation methods on the tibial side in anterior cruciate ligament reconstruction: A biomechanical study in porcine tibial bone. *J Orthop Sci* 2006;11:278-82.
18. Bauer LA, Alberti HA, Corotti VG, Franco AP, Stieven Filho E, Cunha LA. Biomechanical analysis of a double fixation method for tendon graft in porcine tibia – Using an interference screw plus staple. *Rev Bras Ortop* 2018;53:564-9.
19. Pinczewski LA, Clingeleffer AJ, Otto DD, Bonar SF, Corry IS. Integration of hamstring tendon graft with bone in reconstruction of the anterior cruciate ligament. *Arthroscopy* 1997;13:641-3.
20. Weiler A, Hoffmann RF, Bail HJ, Rehm O, Südkamp NP. Tendon healing in a bone tunnel. Part II: Histologic analysis after biodegradable interference fit fixation in a model of anterior cruciate ligament reconstruction in sheep. *Arthroscopy J Arthrosc Related Surg* 2002;18:124-35.
21. Rodeo SA, Arnoczky SP, Torzilli PA, Hidaka C, Warren RF. Tendon-healing in a bone tunnel. A biomechanical and histological study in the dog. *J Bone Joint Surg Am* 1993;75:1795-803.
22. Ghodadra NS, Mall NA, Grumet R, Sherman SL, Kirk S, Provencher MT, *et al.* Interval arthrometric comparison of anterior cruciate ligament reconstruction using bone–patellar tendon–bone autograft versus allograft: Do grafts attenuate within the first year postoperatively? *Am J Sports Med* 2012;40:1347-54.

To,  
The Editor

Covering Letter

**Submission of Manuscript for publication**

Dear Sir,

We intend to publish an article entitled

---

in your journal.

On behalf of all the contributors I will act and guarantor and will correspond with the journal from this point onward.

Prior presentation of the data reported in this manuscript:

Organisation

Place

Date

We have done sufficient work in the field to justify authorship for this manuscript.

We hereby transfer, assign, or otherwise convey all copyright ownership, including any and all rights incidental thereto, exclusively to the journal, in the event that such work is published by the journal.

Thank you,

Yours ' s i n c e r e l y ,

Name of corresponding contributor

Signature

**Title of the manuscript:**

Title Page

**Type of manuscript:**

**Running title:**

**Contributors:**

	First name	Middle name initial	Last name	Highest academic degree	Names of departments and institutions (including city and state)	Email addresses
1						
2						
3						
4						
5						
6						



**Corresponding Author:**

Name:

Address:

Phone numbers:

Facsimile numbers:

E-mail address:

**Total number of pages:**

**Total number of tables:**

**Total number of figures:**

**Total number of supplementary files:**

**Word counts:** For abstract:

For the text:

**Acknowledgement:**

**Conflict of interest:**

**Financial Support:**

Contribution details (to be ticked marked as applicable):

C o n t r i b u

---

	Contributor 1	Contributor 2	Contributor 3	Contributor 4	Contributor 5	Contributor 6
Concepts						
Design						
Definition of intellectual content						
Literature search						
Clinical studies						
Experimental studies						
Data acquisition						
Data analysis						
Statistical analysis						
Manuscript preparation						
Manuscript editing						
Manuscript review						
Guarantor						